



Income level and antibiotic misuse: a systematic review and dose–response meta-analysis

Narmeen Mallah^{1,2,3,4} · Nicola Orsini¹ · Adolfo Figueiras^{2,3,4} · Bahi Takkouche^{2,3,4,5} 

Received: 20 April 2021 / Accepted: 15 November 2021 / Published online: 30 November 2021
© The Author(s) 2021

Abstract

Objectives To quantify the association between income and antibiotic misuse including unprescribed use, storage of antibiotics and non-adherence.

Methods We identified pertinent studies through database search, and manual examination of reference lists of selected articles and review reports. We performed a dose–response meta-analysis of income, both continuous and categorical, in relation to antibiotic misuse. Summary odds ratios (ORs) and their 95% confidence intervals (CIs) were estimated under a random-effects random effects model.

Results Fifty-seven studies from 22 countries of different economic class were included. Overall, the data are in agreement with a flat linear association between income standardized to socio-economic indicators and antibiotic misuse (OR per 1 unit increment = 1.00, p -value = 0.954, p -value non-linearity = 0.429). Data were compatible with no association between medium and high income with general antibiotic misuse (OR 1.04; 95% CI 0.89, 1.20 and OR 1.03; 95% CI 0.82, 1.29). Medium income was associated with 19% higher odds of antibiotic storage (OR 1.19; 95% CI 1.07, 1.32) and 18% higher odds of any aspect of antibiotic misuse in African studies (OR 1.18; 95% CI 1.00, 1.39). High income was associated with 51% lower odds of non-adherence to antibiotic treatment (OR 0.49; 95% CI 0.34, 0.60). High income was also associated with 11% higher odds of any antibiotic misuse in upper-middle wealth countries (OR 1.11; 95% CI 1.00, 1.22).

Conclusions The association between income and antibiotic misuse varies by type of misuse and country wellness. Understanding the socioeconomic properties of antibiotic misuse should prove useful in developing related intervention programs and health policies.

Keywords Income · Antibiotics · Misuse · Meta-analysis · Dose–response

Introduction

The misuse of antibiotics is defined as the intake of these drugs without medical advice (self-prescription) or their use when prescribed by the physician but without compliance with the physician's instructions for treatment regimen in terms of timing, dosage and duration [1, 2]. It is a salient problem worldwide, irrespective of the country economy and wealth. Antibiotic misuse has led to antibiotic resistance, a universal public health problem with high socio-economic and clinical burdens. Different systematic reviews and meta-analyses reported the high prevalence of antibiotic misuse. In their study, Morgan et al. reviewed publications from five continents and concluded that the use of antibiotics without prescription is wide-reaching and accounts for 19 to 100% of antibiotic use outside Northern Europe and North America [3]. Gualano et al. also reported that almost half

✉ Bahi Takkouche
bahi.takkouche@usc.es

¹ Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

² Department of Preventive Medicine, University of Santiago de Compostela, Santiago de Compostela, Spain

³ Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBER-ESP), Madrid, Spain

⁴ Health Research Institute of Santiago de Compostela (IDIS), Santiago de Compostela, Spain

⁵ Department of Preventive Medicine, Faculty of Medicine, University of Santiago de Compostela, 15782 Santiago de Compostela, Spain

of the individuals stop taking antibiotics upon improvement [4]. Another review estimated that the mean use of leftover antibiotics worldwide is 29%, and that of compliance with antibiotic therapy is only 62% [5]. A recent meta-analysis of studies from low- and middle- income countries found that the pooled prevalence of non-prescribed use of antibiotics is considerably high (78%) in these countries [6]. Antibiotic misuse is also frequent in high- income countries, including the United States where the prevalence of antibiotic use without prescription is as high as 66% in some instances, and that of storage of antibiotics for future use ranges between 14 and 48% [7].

Antibiotic resistance causes at least 700,000 annual deaths worldwide [8], more than 35,000 in the United States alone [9]. A similar record is registered in Europe [10]. The impact of antibiotics resistance on the economy is also expanding with disturbing figures [11]. By 2050, the annual mortality rate from antibiotic resistance is projected to exceed that of major causes of death like cancer and diabetes [8], and the provoked economic shortfalls will be as large as that of the 2008–2009 global financial crisis [12].

Several determinants of antibiotic misuse have been identified. These are mainly sociodemographic, including female gender, young adults and elderly, low educational level, difficult access to the healthcare system, unaffordability of the cost of physicians visit and accessibility to antibiotics [7, 13, 14].

In 2012, a narrative review report about self-medication with antibiotics in developing countries analysed data of five studies and concluded that middle income is associated with antibiotic misuse [15]. Studies that evaluated the association of income with antibiotic misuse showed divergent results. Some studies reported up to six-fold increased odds of misuse in high- income individuals [16, 17], while other studies did not find any association [18–20], or detected lower odds of misuse [21, 22]. It is also unclear whether the association between income and antibiotic misuse holds at different social classes and in regions with different levels of access to healthcare and in which regulations about antibiotic dispensing might vary. To the best of our knowledge, there is no meta-analysis that evaluates the association of income with antibiotic misuse worldwide.

To address this gap, we aimed in this study to carry out a meta-analysis of the association of income with antibiotic misuse. We present analyses standardized for socioeconomic indicators.

Materials and methods

PRISMA guidelines were followed for the conduct and reporting of this meta-analysis, and the study protocol was registered in the PROSPERO database (ID: *number deleted*

for blinding purposes). The outcome, antibiotic misuse, was defined as the use or purchase of non-prescribed antibiotics to treat oneself or another person, storage of leftover antibiotics, or nonadherence to the physicians' instructions regarding the dosage, timing and treatment duration. Storage of antibiotics facilitates access to them and therefore is the first step towards their use without prescription [23].

Literature search and study selection

Medline, EMBASE, Conference Proceedings Citation Index-Science, the Open Access Theses and Dissertations, and the five regional bibliographic databases of the World Health Organization (WHO) were searched since their inception until January 2021. The following search syntax was applied in Medline: (Socioeconomic Factors OR income) AND (antibiotic*) AND ((drug storage [MeSH]) OR (compliance) OR (adherence) OR (Nonprescription Drugs/administration & dosage* [MeSH]) OR (misuse) OR (irrational use) OR (left-over)) and adapted for the other databases. We complemented our search by using free text words as follows: antibiotics AND (misuse OR "unprescribed use" OR leftover OR "adherence to treatment") AND (income OR "socioeconomic status" OR "socioeconomic level"). The reference lists of related reviews [3–7, 13–15, 24, 25] and those of included studies were manually checked to supplement database searches. The search was carried out without any language or date restrictions.

Studies that met the following criteria were included: (1) reporting at least two levels of income with defined boundaries as an exposure, and (2) providing Odds Ratio (OR) or Risk Ratio (RR) and their 95% Confidence Interval (CI) as a measurement of the association of income and misuse of antibiotics by the general population, or sufficient data for their calculation.

Data extraction and synthesis

From each included study, we extracted: (1) *general study characteristics*: author's last name and year of publication, study period, participants characteristics (age and gender), and country where the study took place, (2) *exposure*: levels of monthly income, (3) *measures of association*: for each income level: number of subjects who practiced antibiotic misuse, total sample size, adjusted ORs and 95% CIs, and restriction, adjustment, or matching variables. When adjusted ORs were not provided, the crude estimates were registered, and (4) *Type of antibiotic misuse*: use without prescription, non-adherence, and storage of antibiotic leftover. When data were provided for more than one type of antibiotic misuse, we extracted the data of all types of misuse. When the number of events of antibiotic misuse per income level was not available, we contacted the authors to

request this information, but no reply was received [26–28]. We then deemed the number of events missing for those studies. We also inquired about the reference group used in a sub-analysis of one study [29], but due to lack of answer, we did not consider that subgroup.

We standardized the income to country-specific socioeconomic indicators using two approaches. In the first approach, income was standardized to gross domestic product (GDP) *per capita* based on purchasing power parity (PPP) [30]. PPP is a currency conversion rate that is used to equalise the purchasing power of different monetary units. It allows to compare standards of living and economic productivity between countries [31]. In the second approach, the income level was standardized to the adjusted net national income *per capita*, expressed in US dollars [30]. The historical country-specific values of PPP, GDP *per capita* based on PPP, and adjusted net national income *per capita* were extracted from their specific portals in the World Bank [31–33].

Besides data reported in the studies, the classification of countries by economy [34], geographic distribution [35], and literacy rate [36] was obtained.

Statistical analysis

Studies included in this meta-analysis presented income categorized into 2 to 6 levels, with an average of 3 levels. As an estimate of the dose, we used the midpoint assigned to an estimated contrast given the upper and lower boundaries of the income.

We carried out dose–response meta-analysis of income standardized to: (1) gross domestic product (GDP) *per capita* based on (PPP) and (2) adjusted net national income *per capita*.

The dose–response meta-analysis was performed using a one-stage mixed-effects model taking into account heterogeneity across studies [37, 38].

We first used a linear function to estimate a summary OR of antibiotic misuse associated with an increase of 1 unit in income. We next flexibly modelled income using restricted cubic splines with 3 knots fixed at 10th, 50th and 90th percentiles of its distribution. Tests of hypothesis about the regression coefficients of the dose–response model were conducted using a large sample Wald-type test. To facilitate tabular presentation of the summary odds ratios, we further categorized income into tertiles using the lowest as referent.

We stratified the dose–response analysis by type of antibiotic misuse (unprescribed use, storage of leftover, non-adherence); WHO geographic classification, country economy (low wealth, lower–middle wealth, upper–middle wealth and high-wealth); literacy rate ($\geq 90\%$, $< 90\%$); exposure ascertainment (use of pretested or validated questionnaire; untested questionnaire or not reported); comparability

(control for age, sex, educational level and household size; incomplete control); and publication year (≤ 2015 , > 2015). In 2015, WHO published the global action plan to combat the problem of antibiotic resistance [39].

Quality appraisal

As all studies retrieved were eventually of cross-sectional nature, we appraised the quality of the studies using the Newcastle–Ottawa Scale for cross-sectional studies [40]. One point was given for the fulfilment of each of the following criteria: (1) well-defined target population; (2) reported response rate; (3) well described and appropriate statistical analysis; (4) justified sample size; (5) studies adjusted, matched or restricted for age, sex, educational level and household size; (6) use of previously tested or validated questionnaire; and (7) outcome ascertainment carried out using external assessment in addition to self-reporting. When information on a specific criterion was not given, it was graded with 0 point. The grades across items were then summed to obtain a quality score of a maximum of seven points. Two epidemiologists (NM and AF) carried out the quality assessment, and disagreements were resolved by referring to a third epidemiologist (BT).

Publication bias


Publication bias was checked visually using funnel plot and formally through Egger's test [41], and the trim-and-filltrim and fill method [42].

Results

Literature search and study

Figure 1 represents the flow diagram of the selection of studies about income level and misuse of antibiotics. One thousand four hundred fifty-three publications were identified from the literature search, out of which 314 were selected for full-text review (Fig. 1). Fifty-one studies published between 2001 and 2021 met our inclusion criteria (Table 1). Five studies provided data for several types of misuse [20, 27, 28, 43, 44]. We treated each type of misuse as a separate study, making a total of 57 studies introduced in the dose–response analysis. All studies were of cross-sectional design. They involved a total population of 51,008 individuals from 22 countries and 18,094 events of antibiotic misuse. Forty-nine studies were published in English, one in Spanish [45] and one in Croatian [46].

Fig. 1 Flow diagram of the selection of studies about income level and misuse of antibiotics



Income level and antibiotic misuse: continuous analysis

Overall, the data from these 57 studies were compatible with a flat linear association between income standardized to GDP per capita based on PPP and antibiotic misuse (OR 1.00; p -value = 0.954, p -value non-linearity = 0.452). Similar results were obtained for the association of income standardized to adjusted net national income per capita and antibiotic misuse (OR 1.00; p -value = 0.940).

As a graphical presentation of the trend, Fig. 2 shows the estimated summary odds ratio of antibiotic misuse conferred by income standardized to GDP per capita based on PPP.

Income level and antibiotic misuse: categorical and stratified analysis

In the categorical approach of income standardized to GDP per capita based on PPP, overall, as compared to low (1st tertile), no association between income and general antibiotic misuse was observed: medium income (2nd tertile): OR 1.04; 95% CI 0.89, 1.20, and high income (3rd tertile): OR 1.03; 95% CI 0.82, 1.29 (Table 2).

Stratified analysis revealed that medium income was associated with 19% higher odds of *storage of antibiotics* (OR 1.19; 95% CI 1.07, 1.32),; nonetheless, we did not observe any significant association between high income and this type of misuse (OR 1.04; 95% CI 0.92, 1.17). It is noteworthy to mention that *storage of antibiotics* was

Table 1 General characteristics of studies included the dose–response meta-analysis of income level and antibiotic misuse

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Moktan 2021 [63]	India	Attendants of public hospital	18–90	M: 309 F: 195	Use without prescription	37.50	137	41	Reference category 1.10 (0.68–1.77) 1.58 (0.95–2.63) 1.31 (0.67–2.56)	Age, gender, educational level, marital status, public and private clinics, frequency of doctors' consultation, family/friend influence (other family members self-medicating with antibiotics), symptoms (minor illness)
Bulabula 2020 [26]	South Africa	Pregnant women attending public hospital	Mean (SD): 29 (6.1)	F: 301	Use without prescription	49.50	–	–	Reference category 5.40 (0.90–29.90) 4.10 (0.80–19.40) 6.40 (1.20–35.20)	Age, gender, educational level, residential location, knowledge about antibiotics, attitudes towards antibiotics
Chen 2020 [43]	Mali	Medical university students	Mean (SD) 21.3 (2.4)	M: 310 F: 136	Storage of antibiotics	82.95	290	168	Reference category 1.51 (0.96–2.38) 1.31 (0.67–2.56)	Age
					Use without prescription	506.50 1181.60 82.95	114 42 290	77 27 73	Reference category 1.01 (0.62–1.67) 2.46 (1.27–4.77) 1.05 (0.59–1.87)	
Elmahi 2020 [64]	Sudan	General population	≥ 18	M: 130 F: 116	Use without prescription	506.50 1181.60 49.50 149.50	114 42 182 64	29 19 110 38	Reference category	Age, pregnancy, current antibiotic use

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Mallah 2020 [59]	Lebanon	Children's caregivers	≥ 18	M: 276 F: 1092	Any misuse practice	249.50	21	2	Reference category	Age, sex, educational level, area of residence, alcohol consumption, access to medical care facilities, and frequency of telephone medical consultation
						999.50	260	34	1.43 (0.32–6.41)	
						2000.00	223	17	0.78 (0.17–3.65)	
						3000.50	808	36	0.44 (0.10–1.98)	
Nusair 2020 [65]	Jordan	General population	0 to > 65	M: 674 F: 1169	Use without prescription	88.75	175	61	Reference category	Past month antibiotic use
						266.61	659	253	1.16 (0.82–1.65)	
						444.11	1042	458	1.47 (1.05–2.05)	
Rathish 2020 [18]	Sri Lanka	General population	Mean (SD): 36 (21)	M: 181 F: 203	Use without prescription	150.00	267	263	Reference category	NA
						450.00	117	111	2.15 (0.37–12.54)	
Xu 2020 [28]	China	Children's caregivers	Parents with children < 13 years old	M: 1344 F: 4935	Use without prescription	377.50	–	–	Reference category	Age, gender, educational level, medical background, residential location
						1132.58	–	–	0.76 (0.57–1.03)	
						1887.58	–	–	0.81 (0.54–1.21)	
					Storage of antibiotics	377.50	–	–	Reference category	
						1132.58	–	–	1.03 (0.91–1.17)	
						1887.58	–	–	1.16 (0.99–1.36)	
Ateshim 2019 [66]	Eritrea	General population	Median (IQR): 37 (24)	M: 238 F: 339	Use without prescription	0.00	291	–	Reference category	Age, gender, educational level, marital status, occupational status, knowledge about antibiotics, attitudes towards antibiotics
						32.53	92	–	0.92 (0.54–1.56)	
						113.78	136	–	1.22 (0.78–1.19)	
						211.28	58	–	1.43 (0.75–2.73)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Benamer 2019 [67]	Saudi Arabia	University students	Mean (SD): 20.96 (0.148)	M: 166 F: 69	Use without prescription	133.37	164	95	Reference category	Age, gender, educational level, marital status, speciality (medical vs non-medical), residential location, health insurance
						667.50	50	26	0.79 (0.42–1.49)	
						1468.63	18	14	2.54 (0.80–8.06)	
Bogale 2019 [19]	Ethiopia	General population	18 to > 60	M: 246 F: 349	Use without prescription	10.75	–	46	Reference category	Age, gender, educational level, marital status, residential location, occupational status, healthcare profession
						32.27	–	74	2.55 (1.18–5.50)	
						64.52	–	42	1.08 (0.47–2.46)	
						107.52	–	92	1.42 (0.62–3.25)	
Mate 2019 [44]	Mozambique	General population	Median (IQR): 33 (IQR: 25–47)	M: 294 F: 797	Use without prescription	21.24	528	108	Reference category	Age
						63.75	224	45	0.98 (0.66–1.44)	
						127.51	183	40	1.09 (0.72–1.64)	
					Incomplete course of treatment	212.51	117	26	1.11 (0.68–1.80)	
						21.24	506	150	Reference category	
						63.75	215	68	1.10 (0.78–1.55)	
Mukattash 2019 [68]	Jordan	Children's caregivers	20 to ≥ 50	M: 134 F: 712	Use without prescription	127.51	175	60	1.24 (0.86–1.79)	
						212.51	114	21	0.54 (0.32–0.89)	
						352.50	94	41	Reference category	Age
						1058.21	325	141	0.99 (0.62–1.57)	
						1763.21	427	150	0.70 (0.44–1.10)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Sun 2019 [69]	China	Children's caregivers	Parents with children < 13 years old	M: 2243 F: 7283	Storage of antibiotics	230.50	2102	874	Reference category	Age, gender of the parents, gender of the child, educational level, socioeconomic characteristics (residential location and GDP per capita), health insurance, specialty (medical vs non-medical)
						615.50	2889	1434	1.22 (1.08–1.38)	
						1154.00	2749	1355	1.17 (1.02–1.33)	
						1923.00	1786	917	1.36 (1.16–1.60)	
Hu 2018 [70]	China	Medical university students	Mean (SD): 22 (1.5)	M: 661 F: 1158	Use without prescription	768.50	1565	59	Reference category	Age, gender, educational level, parents' educational level, parents medical background, residential location, knowledge-attitudes-and practice score, center of recruitment
						2306.50	254	18	1.95 (1.13–3.36)	
Tong 2018 [71]	China	Attendants of primary care clinics	< 45 to > 60	M: 340 F: 374	Noncompliance	153.20	162	150	Reference category	Age, gender, educational level, residential location, occupation, employment status, knowledge about antibiotics
						344.78	180	163	0.72 (0.33–1.57)	
						651.18	187	158	0.40 (0.20–0.82)	
						880.98	185	150	0.33 (0.16–0.66)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Peng 2018 [20]	China	University students	Guizhou Mean (SD): 21.3 (2.1) Zhejiang Mean (SD): 19.7 (2.6)	M: 2035 F: 1960	Use without prescription	230.92	–	–	Reference category	Age, socioeconomic characteristics (GDP per capita and residential location)
						1001.00	–	–	0.65 (0.39–1.09)	
						2079.08	–	–	0.66 (0.33–1.31)	
					Storage of antibiotics	230.92	–	–	Reference category	
						1001.00	–	–	1.30 (1.10–1.53)	
						2079.08	–	–	1.14 (0.90–1.43)	
Redzick 2018 [46]	Croatia	Attendants of primary care clinics	–	M: 142 F: 402	Use without prescription	230.92	–	–	Reference category	
						1001.00	–	–	1.14 (0.90–1.44)	
						2079.08	–	–	1.05 (0.76–1.46)	
						84.62	88	5	Reference category	Age
						226.12	55	13	5.14 (1.72–15.38)	
						339.32	97	4	0.71 (0.19–2.75)	
Wang 2018 [27]	China	University students	Mean (SD): 20.7 (2.7)	M: 5515 F: 5677	Storage of antibiotics	452.52	100	15	2.93 (1.02–8.42)	
						594.02	199	25	2.39 (0.88–6.45)	
						230.92	3417	–	Reference category	Age, gender, educational level, parents' educational level, parents' medical background, residential location, speciality (medical vs non-medical)
						1001.00	5823	–	1.15 (1.04–1.27)	
						2310.08	1435	–	1.02 (0.88–1.19)	
						3850.08	517	–	1.00 (0.81–1.23)	
Abdelrahman 2017 [60]	Saudi Arabia	General population	<18 to >65	M: 735 F: 293	Use without prescription	230.92	3417	–	Reference category	
						1001.00	5823	–	0.89 (0.67–1.19)	
						2310.08	1435	–	1.13 (0.75–1.71)	
						3850.08	517	–	0.93 (0.53–1.63)	
						200.12	368	112	Reference category	Age
						867.62	146	60	1.59 (1.07–2.37)	
						2002.50	198	72	1.31 (0.91–1.88)	
						3337.63	316	146	1.96 (1.43–2.69)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Albawani 2017 [72]	Yemen	Attendants of pharmacies	Mean (SD): 28.6 (7.7)	M: 204 F: 159	Use without prescription	116.80	268	229	Reference category	Age
						352.80	51	46	1.57 (0.59–4.19)	
Erku 2017 [73]	Ethiopia	General population	Mean (SD): 33.19 (10.82)	M: 163 F: 487	Any misuse practice	581.90	44	41	2.33 (0.69–7.89)	
						50.00	331	282	Reference category	Age, gender, educational level, marital status, employment
						125.50	201	170	0.95 (0.58–1.55)	
Gebrekirstos 2017 [74]	Ethiopia	Attendants of pharmacies	Median (IQR): 30 (16)	M: 473 F: 307	Use without prescription	175.50	118	83	0.41 (0.25–0.68)	size, frequency of visiting health care institutions, satisfaction about healthcare service
						3.26	130	76	1.67 (1.13–2.48)	
						13.00	92	41	0.96 (0.61–1.50)	Age, gender, educational status, marital status, employment
						26.00	81	32	0.78 (0.48–1.26)	
Gillani 2017 [75]	Pakistan	Non-medical university students	Mean (SD): 23.0 (3.4)	M:352 F:375	Use without prescription	39.02	477	218	Reference category	status, household size, residential location, type of illness, healthcare insurance, previous experience with antibiotics, access to health-care
						75.00	245	110	Reference category	Age, specialty (non-medical)
						225.00	180	80	0.98 (0.67–1.45)	
						400.00	136	54	0.81 (0.53–1.24)	
						600.01	166	82	1.20 (0.81–1.78)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Hassali 2017 [76]	Malaysia	General population	Mean (SD): 28.7 (7.4)	M: 171 F: 229	Any misuse practice	124.88	231	82	Reference category	Age, gender, educational level, marital status, race, healthcare related occupation, employment status, health insurance
						499.88	94	29	0.51 (0.27–0.98)	
						1000.00	47	13	0.40 (0.16–0.78)	
						1500.13	28	7	0.42 (0.13–1.34)	
Jamhour 2017 [29]	Lebanon	General population	> 18	M: 182 F: 218	Use without prescription	499.50	88	36	Reference category	Age, gender, educational level, specialty (unrelated to health care)
						1500.00	97	54	1.81 (1.01–3.25)	
Kajeguka 2017 [77]	Tanzania	General population	Mean (SD): 35.4 (13.4)	M: 144 F: 156	Use without prescription	49.50	162	70	2.82 (0.47–16.68)	Age, gender, educational level, marital status, employment status, self-treated condition
						300.50	102	74	1.02 (0.22–4.76)	
						700.50	36	23	Reference category	
Kurniawan 2017 [78]	Indonesia	Attendants of primary care clinics	Median (IQR): 45 (18–49)	M: 137 F: 263	Use without prescription	87.50	186	146	Reference category	Age, gender, educational level, marital status, employment status, health insurance
						262.50	54	34	0.52 (0.24, 1.12)	
Núñez 2017 [79]	Perú	University students	Mean: 19.82	M: 492 F: 508	Use without prescription	462.00	321	204	Reference category	Age
						1386.62	322	211	1.09 (0.79–1.51)	
						2772.62	178	119	1.16 (0.79–1.70)	
						4620.62	179	120	1.17 (0.79–1.72)	
Senadheera 2017 [80]	Sri Lanka	General population	≥ 18	M: 190 F: 174	Use without prescription	87.50	292	15	Reference category	Age, gender, educational level, employment status, health insurance, household size, receiving medical treatment in the last three months, knowledge of antibiotic name
						262.51	288	26	1.83 (0.95–3.54)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Torres 2017 [45]	Ecuador	General population	18–64	M: 97 F: 110	Use without prescription	349.50	200	98	Reference category	Age
						1100.00	132	68	1.11 (0.71–1.72)	
						1775.00	36	14	0.66 (0.32–1.37)	
Aleem 2016 [21]	Saudi Arabia	Children's caregivers	<25 to ≥55	M: 249 F: 382	Use without prescription	2250.50	8	2	0.35 (0.07–1.76)	Age, gender, educational level, household size
						667.50	91	17	Reference category	
						2002.63	519	54	0.50 (0.26, 0.95)	
Bilal 2016 [81]	Pakistan	Attendants of public hospital	Mean (SD): 48.6 (4.4)	M: 263 F: 137	Use without prescription	35.00	180	172	Reference category	Age, residential location, specialty (non-medical related participants)
						105.00	73	62	0.26 (0.10–0.68)	
						210.00	49	36	0.13 (0.05–0.33)	
						415.00	36	29	0.19 (0.06–0.57)	
						685.01	62	26	0.03 (0.01–0.08)	
Zhu 2016 [82]	China	University students	18–45 (IQR: 21–22)	M: 369 F: 291	Use without prescription	40.00	45	28	Reference category	Age, gender, educational level, major, healthcare insurance, residential location
						120.08	423	192	0.50 (0.27–0.95)	
						240.08	173	83	0.56 (0.29–1.10)	
						400.08	19	13	1.32 (0.42–4.11)	
Ding 2015 [83]	China	Children's caregivers	≤29 to ≥50	M: 70 F: 652	Noncompliance	67.08	78	15	Reference category	Age, access to healthcare (number of clinics)
						268.33	384	111	1.71 (0.93–3.13)	
						536.66	260	76	1.73 (0.93–3.24)	
Gebeyehe 2015 [84]	Ethiopia	General population	Mean (SD): <i>Urban</i> 34.1 (12.9) <i>Rural</i> 34.5 (11.5)	M: 263 F: 819	Any misuse practice	25.47	108	30	Reference category	Age, gender, educational level, marital status, employment status, residential location, household size
						76.50	177	59	1.30 (0.77–2.20)	Level of healthcare service satisfaction, knowledge on antibiotics use
						127.53	77	26	1.33 (0.70–2.50)	
						178.53	19	3	0.49 (0.13–1.79)	
						229.53	7	2	1.04 (0.19–5.65)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Yousif 2015 [85]	Saudi Arabia	General population	≥ 18	M: 228 F: 172	Use without prescription	1335.00 4005.13	219 172	173 142	Reference category 0.80 (0.50–1.30)	Age, gender, educational level, marital status, employment status, residential location
Cheaito 2014 [86]	Lebanon	Attendants of pharmacies	Mean (SD): 38.24 (13.7)	M: 143 F: 176	Use without prescription	1000.00 3000.00	278 40	117 17	Reference category 1.02 (0.52–1.99)	Age, gender, educational level, marital status, employment status, health insurance, having a reference doctor and frequency of consultation
Etcha 2014 [87]	Ethiopia	University students	Mean (SD): 21 (2.06)	M: 267 F: 140	Use without prescription	6.28 18.92 31.56	159 160 88	42 38 32	Reference category 0.87 (0.52–1.44) 1.59 (0.91–2.79)	Age, gender, university year, religion, residential location
Hu 2014 [22]	Australia	General population	Mean (SD): 33 (8.2) Range: 14–63	M: 170 F: 258	Storage of antibiotics	1904.13 5712.46	150 278	85 118	Reference category 0.56 (0.38–0.84)	Age, gender, educational level, residential location, employment status, marital status, parental status, language proficiency, main language spoken at home, health insurance
Lv 2014 [88]	China	University students	NA	M:341 F:390	Any misuse practice	41.00 123.08 246.08 410.08	139 447 131 14	58 175 56 5	Reference category 1.14 (0.76–1.71) 1.00 (0.59–1.67) 1.26 (0.39–4.13)	Gender, university year, residential location, major (medical vs non-medical), health insurance

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Mihretie 2014 [89]	Ethiopia	General population	Mean (SD): 37.8 (12.2)	M: 34 F: 17	Use without prescription	13.75	14	9	Reference category	Age
						38.78	10	8	2.22 (0.33–14.80)	
						67.53	10	8	2.22 (0.33–14.80)	
Shah 2014 [90]	Pakistan	University students	Mean (SD): 20.04 (1.74)	M: 253 F: 178	Use without prescription	102.53	14	6	0.42 (0.09–1.91)	
						250.00	115	51	Reference category	Age, specialty (non-medical)
						750.00	139	73	1.39 (0.85–2.28)	
						1250.00	70	38	1.49 (0.82–2.71)	
						1750.01	73	28	0.78 (0.43–1.42)	
Abobtain 2013 [61]	Saudi Arabia	Children's caregivers	<25 to ≥55	M:241 F:369	Use without prescription	667.37	91	17	Reference category	Age, educational level, marital status, household size, number of children <12 years old, healthcare related profession
						2002.50	519	54	0.50 (0.26, 0.95)	Age, gender, major, residential location, healthcare insurance
Pan 2012 [17]	China	University students	Mean (SD): 22.3 (2.6)	M:745 F:555	Use without prescription	38.75	548	215	Reference category	
						116.33	668	352	1.73 (1.37–2.17)	
						232.58	74	46	2.54 (1.54–4.20)	
						387.58	10	8	6.20 (1.30–29.45)	
Widayati 2011 [91]	Indonesia	General population	Median (Range) Prescribed 40.5 (18–69) Self-medicated 43 (18–66)	M: 309 F: 250	Use without prescription	74.50	41	19	Reference category	Age, gender, educational level, marital status, household size, employment status, healthcare insurance
						224.50	24	15	1.93 (0.69–5.40)	
						550.00	5	1	0.29 (0.03–2.82)	
						1050.50	4	2	1.16 (0.15–9.03)	

Table 1 (continued)

Author, Year	Country	Setting	Age (Years)	Sex	Outcome	Mean Income (USD)	Total N/level	Outcome/level	OR point estimate	Adjustment, restriction or matching variables
Ilhan 2009 [16]	Turkey	Attendants of primary care clinics	Mean (SD) 39.5 (15.2)	M:1652 F:1044	Use without prescription	157.43	272	46	Reference category 0.96 (0.67–1.39) 1.32 (0.89–1.97) 1.73 (1.11–2.70) 1.55 (1.02–2.36)	Age, gender, educational level, marital status, employment status, household size, health-care insurance (social security), perceived health status, presence of chronic diseases
Hadi 2008 [92]	Indonesia	Attendants of primary care clinics	Median (range) 31 (0–87)	M: 1147 F: 1849	Use without prescription	13.50 40.50	192 274	30 42	Reference category 0.98 (0.59, 1.63)	Age, gender, educational level, residential location, ethnicity, household size, healthcare insurance
Al-Azzam 2007 [93]	Jordan	General population	≥ 17 to > 60	M:1040 F:1093	Use without prescription	88.75	606	204	Reference category 1.48 (1.18–1.85) 1.36 (1.09–1.69)	NA
Sawair 2007 [94]	Jordan	Attendants of primary care clinics	≤ 16 to > 65	M: 220 F: 257	Use without prescription	266.61 444.11 139.30	721 806 140	309 329 46	Reference category 1.94 (1.18–3.21) 1.35 (0.85–2.14)	Age, gender, educational level, marital status, employment status, healthcare insurance, smoking habits, self-reported health status, chronic comorbidities
Awad 2005 [95]	Sudan	General population	≤ 20 to > 60	M: 790 F: 960	Use without prescription	19.25 67.40 125.15	– – –	– – –	Reference category 0.78 (0.59–1.00) 0.61 (0.42–0.87)	Age, gender, educational level




Fig. 2 Trend of the association of income level standardized to GDP per capita based on PPP and antibiotic misuse. Solid line represents the linear trend. Long-dashed line represents the non-linear restricted cubic spline approach. Short-dashed lines represents 95% CI

evaluated in five studies carried out in China [20, 27, 28, 43, 44] and in a sixth study that was undertaken in Australia but involved Chinese immigrants [22]. High income was associated with 51% lower odds of *non-adherence to antibiotics* treatment (OR 0.49, 95% CI 0.34, 0.70) (Table 2). When restricting the analysis to low-wealth countries, high-income individuals were at 11% higher odds of antibiotic misuse than those with low income in upper-middle wealth countries (OR 1.11; 95% CI 1.00, 1.22) (Table 2). Our findings also suggested an association between medium-income level and antibiotic misuse in African countries (OR 1.18; 95% CI 1.00, 1.39) (Table 2). After 2015, the odds of misuse of antibiotics in medium-income individuals increased when compared with studies undertaken until 2015 (OR_{until 2015} 0.95; 95% CI 0.75, 1.20 and OR_{after 2015} 1.12; 95% CI 0.99, 1.26). Similar findings were obtained for high-income individuals (OR_{until 2015} 0.91; 95% CI 0.62, 1.35 and OR_{after 2015} 1.15; 95% CI 0.93, 1.41) (Table 2).

Table 2 Meta-analysis of the association of income level represented as units of GDP per capita based on PPP with antibiotic misuse

	Number of studies	Medium income OR (95% CI)	High income OR (95% CI)
All studies	57	1.04 (0.89, 1.20)	1.03 (0.82, 1.29)
Type of misuse			
Use without prescription	43	1.06 (0.87, 1.28)	1.07 (0.84, 1.37)
Storage of antibiotics	6	1.19 (1.07, 1.32)	1.04 (0.92, 1.17)
Non-adherence	3	1.10 (0.89, 1.35)	0.49 (0.34, 0.70)
Country economy			
Low	16	1.02 (0.83, 1.24)	0.90 (0.59, 1.37)
Lower-middle	11	1.14 (0.73, 1.80)	0.92 (0.46, 1.84)
Upper-middle	25	1.17 (0.91, 1.49)	1.11 (1.00, 1.22)
High	5	0.90 (0.44, 1.85)	1.04 (0.33, 3.28)
WHO Region			
African	14	1.18 (1.00, 1.39)	0.96 (0.67, 1.38)
Eastern Mediterranean	17	0.92 (0.65, 1.32)	0.95 (0.58, 1.57)
South-East Asian	6	1.11 (0.62, 2.00)	1.53 (0.81, 2.92)
Western Pacific	16	0.99 (0.82, 1.20)	1.05 (0.92, 1.19)
Survey year			
Until 2015	29	0.95 (0.75, 1.20)	0.91 (0.62, 1.35)
After 2015	28	1.12 (0.99, 1.26)	1.15 (0.93, 1.41)
Literacy rate			
< 90%	20	1.03 (0.82, 1.29)	1.02 (0.68, 1.54)
≥ 90%	37	1.09 (0.93, 1.28)	1.02 (0.84, 1.23)
Pre-tested or validated questionnaire			
No	10	1.02 (0.51, 2.06)	0.90 (0.34, 2.36)
Yes	47	1.06 (0.91, 1.24)	1.04 (0.85, 1.27)
Adjustment			
Incomplete	47	1.09 (0.95, 1.24)	1.05 (0.84, 1.31)
Complete	10	0.90 (0.71, 1.15)	0.60 (0.30, 1.23)
Quality Score			
Lower quality (≤ 3 points)	24	0.99 (0.75, 1.31)	1.09 (0.72, 1.66)
Higher quality (> 3 points)	33	1.04 (0.86, 1.25)	1.03 (0.81, 1.31)

No meaningful difference in the odds of antibiotic misuse by medium- and high- income individuals was observed when countries were grouped according to literacy rate (Table 2).

The categorical approach of income standardized to net national income per capita showed similar results to that of income standardized to GDP per capita based on PPP (data not shown).

Methodological characteristics of the studies

Restricting the analysis to those studies that used pretested or validated questionnaires did not yield any substantial modification in the pooled OR estimates (OR_{medium} 1.06; 95% CI 0.91, 1.24 and OR_{high} 1.04; 95% CI 0.85, 1.27) (Table 2).

Studies that incompletely controlled for sex, age, educational level and household size showed higher pooled estimates than those with complete control of those variables in medium income ($OR_{\text{incomplete}}$ 1.09; 95% CI 0.95, 1.24 and OR_{complete} 0.90; 95% CI 0.71, 1.15) and in high income ($OR_{\text{incomplete}}$ 1.05; 95% CI 0.84, 1.31 and OR_{complete} 0.60; 95% CI 0.30, 1.23) (Table 2).

No notable difference was observed between pooled estimates from studies with lower-quality (≤ 3 points) and those from studies with higher-quality score (> 3 points) (Table 2).

Publication bias

The funnel plot (Fig. 3) and Egger's test of the null hypothesis (p -value = 0.39) did not suggest evidence of publication bias. These findings were further confirmed by




Fig. 3 Funnel plot of studies about income and antibiotic misuse

the Trim-and-Fill analysis that did not yield to the addition of any study.

Discussion

Antibiotic resistance is an internationally growing multifaceted emergency that has been exacerbated by antibiotic misuse and has left devastating impact at the clinical, health and socio-economic levels. If not controlled, antibiotic resistance will convert into the major cause of death in 2050 [8].

To the best of our knowledge, this is the first meta-analysis that assesses the dose–response association between income level and misuse of antibiotics. Our results agree well with the hypothesis of no association between income level and misuse of antibiotics. Subgroup analyses reveal a dose–response association of medium- and high- income levels with specific types of antibiotic misuse, i.e., storage of drug leftover and non-adherence, country wealth, geographic region and study period.

Our primary findings suggest that the odds of misuse of antibiotics do not differ between poor and wealthy people. This is in line with the fact that both low- and high- income individuals tend to self-medicate. On the one hand, under constrained financial resources, especially in less developed economies where access to health facilities is limited, self-medication is the only available option of healthcare [47]. By self-medicating, individuals with low income avoid expenses of medical consultation and subsequent lab tests. Low- income households report forgone care more often than those with high- income level [48]. They often cut back basic needs and take less medication than prescribed, due to cost [49, 50], explaining therefore the observed higher likelihood of adherence to treatment by high- income than by low- income individuals. On the other hand, people with high- income level tend to medicate themselves as they have easier access to sources of information including internet to seek health information [51], can afford purchasing non-reimbursed medicines, and have more social support that increases their access to unprescribed medicines including through sharing with families and friends [52].

Our dose–response meta-analysis also showed that medium- income individuals have higher odds of storing antibiotic leftover than those with low income. This could be related to higher financial affordability by medium-income medium income individuals to purchase and store antibiotics. Our results also show a higher likelihood of misuse by high-income individuals in upper–middle wealth countries. Consistent with our findings, an earlier report about the economy of self-medication in general, indicated that the demand for self-medication declines with rising the income level of high- income individuals, but increases with

increasing the income of low-income individuals, resulting in a null pooled effect between income and self-medication [47].

We also reported that medium- income individuals in Africa have higher chances of antibiotic misuse, probably due to the poor enforcement of antibiotic dispensing regulations in those regions.

We observed a marginal increase in the odds of misuse of antibiotics by medium- income and high- income individuals after 2015 than before this period. This could be related to two main motives: first, as concluded by WHO in its report *Global Spending on Health*, the expenditure on health is growing faster than economies, leading to a doubling of the out-of-pocket spending and very large differences between high- and low-wealth countries concerning health expenditure [53], second, not all countries have developed and implemented sufficient measures to control the dispensing of antibiotics, and thus people with greater financial resources continued using antibiotics without prescription. A recent review report indicated that more than half of the antibiotics worldwide are dispensed without prescription [54]. Consequently, the WHO placed a new urgent call to control antibiotics resistance crisis on 2019 [55].

The findings of this meta-analysis are unlikely to be affected by publication bias as revealed by the negative result of Egger's test and the trim-and-fill analysis that did not suggest imputation of any additional study.

This meta-analysis suffers from several limitations. All eligible studies were of cross-sectional design, which, theoretically, limited any causal inference. However, income is a relatively stable variable through time and, which mitigates this limitation. Furthermore, only one-fifth of included studies performed a complete control for socio-demographic variables, and higher OR estimates were obtained from studies with incomplete adjustment than in studies with complete adjustment. This reveals that our findings could be overestimated due to incomplete adjustment. Additional studies that control adequately for all potentially related socio-demographic variables are needed to confirm our results. Also, one-sixth of studies did not employ a pretested or validated questionnaire to ascertain the exposure and the outcome. However, this was unlikely to affect our results as constraining the analysis to the remaining studies did not introduce any change in the overall effect.

Our analysis was based on random-effect models to account for heterogeneity between studies [56–58]. Heterogeneity was expected in our study due to difference in the defined levels of income, period of antibiotic use (for example, use in the past month [59], past 3 months [60] and past year [61]), and settings. Experts in meta-analysis emphasize that heterogeneity is the expectation in any meta-analysis rather than the exception [62] and that no amount of heterogeneity is considered unacceptable as long as the

inclusion criteria are clearly defined and the data are correctly analysed [56].

Understanding the socioeconomic properties of antibiotic misuse is crucial to develop related intervention programs and health policies, yet addition of high-quality studies that control for socio-demographic and socio-economic indicators are needed to confirm our findings.

Acknowledgements Mrs Narmeen Mallah received a Grant for her internship at Karolinska Institutet from Erasmus+ KA103 Erasmus European Mobility Program. The authors would like to thank to Mr. Luís Cea and Dr. Sami Ashour for their help with economic concepts.

Author contributions NM and BT conceived the research idea, carried out the literature review and extracted the data. AF participated in quality assessment of retrieved studies. NM carried out data analysis and interpretation and designed and wrote the manuscript. BT and NO supervised data analyses. All authors reviewed and revised the manuscript and approved it for publication.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Data availability The data generated and analyzed in the meta-analysis are included in the article. The data are available by accessing the cited references.

Declarations

Conflict of interest The authors declare no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Centers for Disease Control and Prevention: Antibiotic use questions and answers. https://www.cdc.gov/antibiotic-use/community/about/should-know.html#anchor_1572453021219 (2019). Accessed 21 Apr 2021
- United Nations Office on Drugs and Crime: The non-medical use of prescription drugs: Policy direction issues. <https://www.unodc.org/documents/drug-prevention-and-treatment/nonmedical-use-prescription-drugs.pdf> (2011). Accessed 20 Aug 2021
- Morgan, D.J., Okeke, I.N., Laxminarayan, R., Perencevich, E.N., Weisenberg, S.: Non-prescription antimicrobial use worldwide: a systematic review. *Lancet Infect. Dis.* **11**(9), 692–701 (2011). [https://doi.org/10.1016/S1473-3099\(11\)70054-8](https://doi.org/10.1016/S1473-3099(11)70054-8)
- Gualano, M.R., Gili, R., Scaioi, G., Bert, F., Siliquini, R.: General population's knowledge and attitudes about antibiotics: a

- systematic review and meta-analysis. *Pharmacoepidemiol. Drug Saf.* **24**(1), 2–10 (2015). <https://doi.org/10.1002/pds.3716>
5. Kardas, P., Devine, S., Golembesky, A., Roberts, C.: A systematic review and meta-analysis of misuse of antibiotic therapies in the community. *Int. J. Antimicrob. Agents* **26**(2), 106–113 (2005). <https://doi.org/10.1016/j.ijantimicag.2005.04.017>
 6. Torres, N.F., Chibi, B., Kuupiel, D., Solomon, V.P., Mashamba-Thompson, T.P., Middleton, L.E.: The use of non-prescribed antibiotics; prevalence estimates in low-and-middle-income countries. A systematic review and meta-analysis. *Arch. Public Health.* **79**(1), 2 (2021). <https://doi.org/10.1186/s13690-020-00517-9>
 7. Grigoryan, L., Germanos, G., Zoorob, R., Juneja, S., Raphael, J.L., Paasche-Orlow, M.K., et al.: Use of antibiotics without a prescription in the U.S. population: a scoping review. *Ann. Intern. Med.* **171**(4), 257–263 (2019). <https://doi.org/10.7326/M19-0505>
 8. Wellcome Trust: Review on antimicrobial resistance. Tackling drug-resistant infections globally: Final report and recommendations. https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf (2016). Accessed 21 Apr 2021
 9. Centers for Disease Control and Prevention: Antibiotic resistance threats in the United States. www.cdc.gov/DrugResistance/Biggest-Threats.html (2019). Accessed 21 Apr 2021
 10. European Center for Disease Prevention and Control & Organization for Economic Co-operation and Development: Antimicrobial resistance tackling the burden in the European Union. A Briefing note for EU/EEA countries. <https://www.ecdc.europa.eu/en/antimicrobial-resistance> (2019). Accessed 21 Apr 2021
 11. Spellberg, B., Blaser, M., Guidos, R.J., Boucher, H.W., Bradley, J.S., Eisenstein, B.I., et al.: Combating antimicrobial resistance: policy recommendations to save lives. *Clin. Infect. Dis.* **52**(Suppl 5), S397–428 (2011). <https://doi.org/10.1093/cid/cir153>
 12. World Bank: Drug-resistant infections: A threat to our economic future. Washington, DC. <https://documents.worldbank.org/en/publication/documents-reports/documentdetail/323311493396993758/final-report> (2017). Accessed 21 Apr 2021
 13. Alhomoud, F., Aljamea, Z., Almahasnah, R., Alkhalifah, K., Basalelah, L., Alhomoud, F.K.: Self-medication and self-prescription with antibiotics in the Middle East-do they really happen? A systematic review of the prevalence, possible reasons, and outcomes. *Int J Infect Dis.* **57**, 3–12 (2017). <https://doi.org/10.1016/j.ijid.2017.01.014>
 14. Torres, N.F., Chibi, B., Middleton, L.E., Solomon, V.P., Mashamba-Thompson, T.P.: Evidence of factors influencing self-medication with antibiotics in low and middle-income countries: a systematic scoping review. *Public Health* **168**, 92–101 (2019). <https://doi.org/10.1016/j.puhe.2018.11.018>
 15. Ocan, M., Obuku, E.A., Bwanga, F., Akena, D., Richard, S., Ogwal-Okeng, J., et al.: Household antimicrobial self-medication: a systematic review and meta-analysis of the burden, risk factors and outcomes in developing countries. *BMC Public Health* **15**, 742 (2015). <https://doi.org/10.1186/s12889-015-2109-3>
 16. Ilhan, M.N., Durukan, E., Ilhan, S.O., Aksakal, F.N., Ozkan, S., Bumin, M.A.: Self-medication with antibiotics: questionnaire survey among primary care center attendants. *Pharmacoepidemiol. Drug Saf.* **18**(12), 1150–1157 (2009). <https://doi.org/10.1002/pds.1829>
 17. Pan, H., Cui, B., Zhang, D., Farrar, J., Law, F., Ba-Thein, W.: Prior knowledge, older age, and higher allowance are risk factors for self-medication with antibiotics among university students in southern China. *PLoS ONE* **7**(7), e41314 (2012). <https://doi.org/10.1371/journal.pone.0041314>
 18. Rathish, D., Wickramasinghe, N.D.: Prevalence, associated factors and reasons for antibiotic self-medication among dwellers in Anuradhapura: a community-based study. *Int. J. Clin. Pharm.* **42**(4), 1139–1144 (2020). <https://doi.org/10.1007/s11096-020-01065-6>
 19. Bogale, A.A., Amhare, A.F., Chang, J., Bogale, H.A., Betaw, S.T., Gebrehiwot, N.T., et al.: Knowledge, attitude, and practice of self-medication with antibiotics among community residents in Addis Ababa, Ethiopia. *Expert Rev. Anti Infect Ther.* **17**(6), 459–466 (2019). <https://doi.org/10.1080/14787210.2019.1620105>
 20. Peng, D., Wang, X., Xu, Y., Sun, C., Zhou, X.: Antibiotic misuse among university students in developed and less developed regions of China: a cross-sectional survey. *Glob. Health Action* **11**(1), 1496973 (2018). <https://doi.org/10.1080/16549716.2018.1496973>
 21. Aleem, M.A., Rahman, M., Ishfaq, M., Mehmood, K., Ahmed, S.S.: Determinants of antibiotics misuse by the parents in children: a survey from Northern Region of Saudi Arabia. *Bangladesh J. Child Health* **40**(2), 64–71 (2016)
 22. Hu, J., Wang, Z.: In-home antibiotic storage among Australian Chinese migrants. *Int. J. Infect. Dis.* **26**, 103–106 (2014). <https://doi.org/10.1016/j.ijid.2014.04.017>
 23. Grigoryan, L., Monnet, D.L., Haaijer-Ruskamp, F.M., Bonten, M.J., Lundborg, S., Verheij, T.J.: Self-medication with antibiotics in Europe: a case for action. *Curr. Drug Saf.* **5**(4), 329–332 (2010). <https://doi.org/10.2174/157488610792246046>
 24. De Sanctis, V., Soliman, A.T., Daar, S., Di Maio, S., Elalaily, R., Fiscina, B., et al.: Prevalence, attitude and practice of self-medication among adolescents and the paradigm of dysmenorrhea self-care management in different countries. *Acta Biomed.* **91**(1), 182–192 (2020). <https://doi.org/10.23750/abm.v91i1.9242>
 25. Xu, R., Mu, T., Wang, G., Shi, J., Wang, X., Ni, X.: Self-medication with antibiotics among university students in LMIC: a systematic review and meta-analysis. *J. Infect. Dev. Countries* **13**(8), 678–689 (2019). <https://doi.org/10.3855/jidc.11359>
 26. Bulabula, A.N.H., Dramowski, A., Mehtar, S.: Antibiotic use in pregnancy: knowledge, attitudes and practices among pregnant women in Cape Town South Africa. *J. Antimicrob. Chemother.* **75**(2), 473–481 (2020). <https://doi.org/10.1093/jac/dkz427>
 27. Wang, X., Lin, L., Xuan, Z., Li, L., Zhou, X.: Keeping antibiotics at home promotes self-medication with antibiotics among Chinese university students. *Int. J. Environ. Res. Public Health* **15**(4), 687 (2018). <https://doi.org/10.3390/ijerph15040687>
 28. Xu, Y., Lu, J., Sun, C., Wang, X., Hu, Y.J., Zhou, X.: A cross-sectional study of antibiotic misuse among Chinese children in developed and less developed provinces. *J. Infect. Dev. Countries* **14**(2), 129–137 (2020). <https://doi.org/10.3855/jidc.11938>
 29. Jamhour, A., El-Kheir, A., Salameh, P., Hanna, P.A., Mansour, H.: Antibiotic knowledge and self-medication practices in a developing country: a cross-sectional study. *Am. J. Infect. Control* **45**(4), 384–388 (2017). <https://doi.org/10.1016/j.ajic.2016.11.026>
 30. The World Bank: Metadata glossary. <https://databank.worldbank.org/metadataglossary/world-development-indicators/series/NY.ADJ.NNTY.CD>. Accessed 1 Mar 2021
 31. The World Bank: Price level ratio of PPP conversion factor (GDP) to market exchange rate. <https://data.worldbank.org/indicator/PA.NUS.PPPC.RF?view=chart>. Accessed 8 Feb 2021
 32. The World Bank: GDP per capita, PPP (current international \$). <https://data.worldbank.org/indicator/NY.GDP.PCAP.PP.CD>. Accessed 8 Feb 2021
 33. The World Bank: Adjusted net national income per capita (current US\$). <https://data.worldbank.org/indicator/NY.ADJ.NNTY.PC.CD>. Accessed 8 Feb 2021
 34. The World Bank: World Bank country and lending groups, country classification. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>. Accessed 24 Feb 2021
 35. World Health Organization: Countries. <https://www.who.int/countries/>. Accessed 18 Feb 2021

36. United Nations Educational Scientific and Cultural Organization: Education and literacy. <http://uis.unesco.org/en/country/lb> (2018). Accessed 21 Apr 2021
37. Orsini, N.: Weighted mixed-effects dose-response models for tables of correlated contrasts. *Stata J.* **21**, 320 (2021)
38. Crippa, A., Discacciati, A., Bottai, M., Spiegelman, D., Orsini, N.: One-stage dose-response meta-analysis for aggregated data. *Stat. Methods Med. Res.* **28**(5), 1579–1596 (2019). <https://doi.org/10.1177/0962280218773122>
39. World Health Organization: Global action plan on antimicrobial resistance. <https://www.who.int/antimicrobial-resistance/publications/global-action-plan/en/>. (2015). Accessed 21 Apr 2021
40. Modesti, P.A., Reboldi, G., Cappuccio, F.P., Agyemang, C., Remuzzi, G., Rapi, S., et al.: Panethnic differences in blood pressure in Europe: a systematic review and meta-analysis. *PLoS ONE* **11**(1), e0147601 (2016). <https://doi.org/10.1371/journal.pone.0147601>
41. Egger, M., Davey Smith, G., Schneider, M., Minder, C.: Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**(7109), 629–634 (1997). <https://doi.org/10.1136/bmj.315.7109.629>
42. Duval, S., Tweedie, R.: Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* **56**(2), 455–463 (2000). <https://doi.org/10.1111/j.0006-341x.2000.00455.x>
43. Chen, J., Sidibi, A.M., Shen, X., Dao, K., Maiga, A., Xie, Y., et al.: Lack of antibiotic knowledge and misuse of antibiotics by medical students in Mali: a cross-sectional study. *Expert Rev. Anti Infect. Ther.* **19**, 1–8 (2020). <https://doi.org/10.1080/14787210.2021.1857731>
44. Mate, I., Come, C.E., Goncalves, M.P., Cliff, J., Gudo, E.S.: Knowledge, attitudes and practices regarding antibiotic use in Maputo City, Mozambique. *PLoS ONE* **14**(8), e0221452 (2019). <https://doi.org/10.1371/journal.pone.0221452>
45. Torres, K.S., Ochoa, A., Encalada, D., Quizhpe, A.: Prevalence of self-medication with antibiotics in the urban parishes of the city of Cuenca, 2016–2017 [Prevalencia de la automedicación con antibióticos en las parroquias urbanas de la ciudad de Cuenca, 2016–2017]. *Archivos Venezolanos de Farmacología y Terapéutica* **36**(4), 130–136 (2017)
46. Redzic, L., Zalihic, A.: Self-medication with antibiotics in family practice in patients and parents [Samomedikacija antibioticima među pacijentima]. *Croat. J. Infect. [Infektološki glasnik]* **38**(3), 69–73 (2018)
47. Chang, F.R., Trivedi, P.K.: Economics of self-medication: theory and evidence. *Health Econ.* **12**(9), 721–739 (2003). <https://doi.org/10.1002/hec.841>
48. Mielck, A., Kiess, R., von dem Knesebeck, O., Stirbu, I., Kunst, A.E.: Association between forgone care and household income among the elderly in five Western European countries—analyses based on survey data from the SHARE-study. *BMC Health Serv. Res.* **9**, 52 (2009). <https://doi.org/10.1186/1472-6963-9-52>
49. Mojtabai, R., Olfson, M.: Medication costs, adherence, and health outcomes among Medicare beneficiaries. *Health Aff. (Millwood)* **22**(4), 220–229 (2003). <https://doi.org/10.1377/hlthaff.22.4.220>
50. Piette, J.D., Heisler, M., Wagner, T.H.: Cost-related medication underuse among chronically ill adults: the treatments people forgo, how often, and who is at risk. *Am. J. Public Health* **94**(10), 1782–1787 (2004). <https://doi.org/10.2105/ajph.94.10.1782>
51. Renahy, E., Parizot, I., Chauvin, P.: Health information seeking on the Internet: a double divide? Results from a representative survey in the Paris metropolitan area, France, 2005–2006. *BMC Public Health* **8**, 69 (2008). <https://doi.org/10.1186/1471-2458-8-69>
52. Vanhaesebrouck, A., Vuillermoz, C., Robert, S., Parizot, I., Chauvin, P.: Who self-medicates? Results from structural equation modeling in the Greater Paris area, France. *PLoS ONE* **13**(12), e0208632 (2018). <https://doi.org/10.1371/journal.pone.0208632>
53. World Health Organization: Global spending on health: a world in transition. https://www.who.int/health_financing/documents/health-expenditure-report-2019/en/. (2019). Accessed 21 April 2021
54. Batista, A.D., Rodrigues, D.A., Figueiras, A., Zapata-Cachafeiro, M., Roque, F., Herdeiro, M.T.: Antibiotic dispensation without a prescription worldwide: a systematic review. *Antibiotics (Basel)* **9**(11), 786 (2020). <https://doi.org/10.3390/antibiotics9110786>
55. Interagency Coordination Group on Antimicrobial Resistance: No time to wait: Securing the future from drug-resistant infections. Report to the secretary-general of the United Nations.: World Health Organization. <https://www.who.int/antimicrobial-resistance/interagency-coordination-group/final-report/en/>. (2019). Accessed 21 Apr 2021
56. Higgins, J.P.: Commentary: heterogeneity in meta-analysis should be expected and appropriately quantified. *Int. J. Epidemiol.* **37**(5), 1158–1160 (2008). <https://doi.org/10.1093/ije/dyn204>
57. National Research Council: Combining Information: Statistical Issues and Opportunities for Research. The National Academies Press, Washington, DC (1992)
58. Murad, M.H., Montori, V.M., Ioannidis, J.P.A., et al.: Fixed-effects and random-effects models. In: Guyatt, G., Rennie, D., Meade, M.O. (eds.) *Users' Guide to the Medical Literature A Manual for Evidence-Based Clinical Practice*, 3rd edn., p. 885. McGraw-Hill, New York (2015)
59. Mallah, N., Badro, D.A., Figueiras, A., Takkouche, B.: Association of knowledge and beliefs with the misuse of antibiotics in parents: a study in Beirut (Lebanon). *PLoS ONE* **15**(7), e0232464 (2020). <https://doi.org/10.1371/journal.pone.0232464>
60. Abdelrahman, T.M., Al Saeed, M.S., Karam, R.A., Alkhthami, A.M., Alswat, O.B., Alzahrani, A.A., et al.: Misuse of antibiotics and antibiotic resistance: a public population-based health survey in al Taif- Saudi Arabia. *WJPMR* **3**(2), 54–62 (2017)
61. Abobotain, A.H., Sheerah, H.A., Alotaibi, F.N., Joury, A.U., Mishiddi, R.M., Siddiqui, A.R., et al.: Socio-demographic determinants of antibiotic misuse in children. A survey from the central region of Saudi Arabia. *Saudi Med. J.* **34**(8), 832–840 (2013)
62. Berlin, J.A.: Invited commentary: benefits of heterogeneity in meta-analysis of data from epidemiologic studies. *Am. J. Epidemiol.* **142**(4), 383–387 (1995). <https://doi.org/10.1093/oxfordjournals.aje.a117645>
63. Moktan, D., Shehnaz, S.I.: Factors driving self-medication with antimicrobials in Karaikal, Puducherry India. *J. Pharmacol. Pharmacother.* **11**, 64–71 (2021). https://doi.org/10.4103/jpp.JPP_21_20
64. Elmahi, O.K.O., Balla, S.A., Khalil, H.A.: Self-medication with antibiotics and its predictors among the population in Khartoum Locality, Khartoum State, Sudan in 2018. *Int. J. Trop. Dis. Health* **41**(4), 17–25 (2020). <https://doi.org/10.9734/IJTDH/2020/v4i430267>
65. Nusair, M.B., Al-Azzam, S., Alhamad, H., Momani, M.Y.: The prevalence and patterns of self-medication with antibiotics in Jordan: a community-based study. *Int. J. Clin. Pract.* **75**, e13665 (2021). <https://doi.org/10.1111/ijcp.13665>
66. Ateshim, Y., Bereket, B., Major, F., Emun, Y., Woldai, B., Pasha, I., et al.: Prevalence of self-medication with antibiotics and associated factors in the community of Asmara, Eritrea: a descriptive cross sectional survey. *BMC Public Health* **19**(1), 726 (2019). <https://doi.org/10.1186/s12889-019-7020-x>
67. Benameur, T., Al-Bohassan, H., Al-Aithan, A., Al-Beladi, A., Al-Ali, H., Al-Omran, H., et al.: Knowledge, attitude, behaviour of the future healthcare professionals towards the self-medication practice with antibiotics. *J. Infect. Dev. Countries* **13**(1), 56–66 (2019). <https://doi.org/10.3855/jidc.10574>

68. Mukattasha, T.L., Alkhatatbeha, M.J., Andrawosa, S., Jaraba, A.S., AbuFarhab, R.K., Nusair, M.B.: Parental self-medication of antibiotics for children in Jordan. *JPHSR* (2019). <https://doi.org/10.1111/jphs.12331>
69. Sun, C., Hu, Y.J., Wang, X., Lu, J., Lin, L., Zhou, X.: Influence of leftover antibiotics on self-medication with antibiotics for children: a cross-sectional study from three Chinese provinces. *BMJ Open* **9**(12), e033679 (2019). <https://doi.org/10.1136/bmjopen-2019-033679>
70. Hu, Y., Wang, X., Tucker, J.D., Little, P., Moore, M., Fukuda, K., et al.: Knowledge, attitude, and practice with respect to antibiotic use among Chinese medical students: a multicentre cross-sectional study. *Int. J. Environ. Res. Public Health* **15**(6), 1165 (2018). <https://doi.org/10.3390/ijerph15061165>
71. Tong, S., Pan, J., Lu, S., Tang, J.: Patient compliance with antimicrobial drugs: a Chinese survey. *Am. J. Infect. Control* **46**(4), e25–e29 (2018). <https://doi.org/10.1016/j.ajic.2018.01.008>
72. Albawani, S.M., Hassan, Y.B., Abd-Aziz, N., Gnanasan, S.: Self-medication with antibiotics in Sana'a City, Yemen. *Trop. J. Pharm. Res.* **16**(5), 1195 (2017). <https://doi.org/10.4314/tjpr.v16i5.30>
73. Erku, D.A., Mekuria, A.B., Belachew, S.A.: Inappropriate use of antibiotics among communities of Gondar town, Ethiopia: a threat to the development of antimicrobial resistance. *Antimicrob. Resist. Infect. Control* **6**, 112 (2017). <https://doi.org/10.1186/s13756-017-0272-2>
74. Gebrekirstos, N.H., Workneh, B.D., Gebregiorgis, Y.S., Misgina, K.H., Weldehaweria, N.B., Weldu, M.G., et al.: Non-prescribed antimicrobial use and associated factors among customers in drug retail outlet in Central Zone of Tigray, Northern Ethiopia: a cross-sectional study. *Antimicrob. Resist. Infect. Control* **6**, 70 (2017). <https://doi.org/10.1186/s13756-017-0227-7>
75. Gillani, A.H., Ji, W., Hussain, W., Imran, A., Chang, J., Yang, C., et al.: Antibiotic self-medication among non-medical university students in Punjab, Pakistan: a cross-sectional survey. *Int. J. Environ. Res. Public Health* **14**(10), 1152 (2017). <https://doi.org/10.3390/ijerph14101152>
76. Hassali, M.A., Arief, M., Saleem, F., Khan, M.U., Ahmad, A., Mariam, W., et al.: Assessment of attitudes and practices of young Malaysian adults about antibiotics use: a cross-sectional study. *Pharm. Pract. (Granada)* **15**(2), 929 (2017). <https://doi.org/10.18549/PharmPract.2017.02.929>
77. Kajeguka, D.C., Moses, E.A.: Self-medication practices and predictors for self-medication with antibiotics and antimalarials among community in Mbeya City, Tanzania. *Tanzan. J. Health Res.* (2017). <https://doi.org/10.4314/thrb.v19i4.6>
78. Kurniawan, K., Posangi, J., Rampengan, N.: Association between public knowledge regarding antibiotics and self-medication with antibiotics in Teling Atas Community Health Center East Indonesia. *Med. J. Indones.* **25**, 62–69 (2017). <https://doi.org/10.13181/mji.v26i1.1589>
79. Nuñez, M., Tresierra-Ayalab, M., Gil-Olivares, F.: Antibiotic self-medication in university students from Trujillo, Peru. *Medicina Universitaria* **18**(73), 205–209 (2017). <https://doi.org/10.1016/j.rmu.2016.10.003>
80. Senadheera, G.P., Sri Ranganathan, S., Gunawardane, N.S., Fernando, G.H., Fernandopulle, B.M.: Practice of self-medication with antibiotics in the Colombo district, Sri Lanka. *Ceylon Med. J.* **62**(1), 70–72 (2017). <https://doi.org/10.4038/cmj.v62i1.8439>
81. Bilal, M., Haseeb, A., Khan, M.H., Arshad, M.H., Ladak, A.A., Niazi, S.K., et al.: Self-medication with antibiotics among people dwelling in rural areas of Sindh. *J. Clin. Diagn. Res.* **10**(5), 08–13 (2016). <https://doi.org/10.7860/JCDR/2016/18294.7730>
82. Zhu, X., Pan, H., Yang, Z., Cui, B., Zhang, D., Ba-Thein, W.: Self-medication practices with antibiotics among Chinese university students. *Public Health* **130**, 78–83 (2016). <https://doi.org/10.1016/j.puhe.2015.04.005>
83. Ding, L., Sun, Q., Sun, W., Du, Y., Li, Y., Bian, X., et al.: Antibiotic use in rural China: a cross-sectional survey of knowledge, attitudes and self-reported practices among caregivers in Shandong province. *BMC Infect. Dis.* **15**, 576 (2015). <https://doi.org/10.1186/s12879-015-1323-z>
84. Gebeyehu, E., Bantie, L., Azage, M.: Inappropriate use of antibiotics and its associated factors among urban and rural communities of Bahir Dar City administration, Northwest Ethiopia. *PLoS ONE* **10**(9), e0138179 (2015). <https://doi.org/10.1371/journal.pone.0138179>
85. Yousif, M.A., Abubaker, I.E.: Prevalence, determinants and practices of self-medication with antibiotics: a population based survey in Taif, Kingdom of Saudi Arabia. *Int. J. Res. Pharm. Sci.* **5**(2), 51–56 (2015)
86. Cheaito, L., Azizi, S., Saleh, N., Salameh, P.: Assessment of self-medication in population buying antibiotics in pharmacies: a pilot study from Beirut and its suburbs. *Int. J. Public Health* **59**(2), 319–327 (2014). <https://doi.org/10.1007/s00038-013-0493-y>
87. Eticha, T., Araya, H., Alemayehu, A., Solomon, G., Ali, D.: Prevalence and predictors of self-medication with antibiotics among Adi-haqi Campus students of Mekelle University, Ethiopia. *IJPSR* **5**(10), 678 (2014)
88. Lv, B., Zhou, Z., Xu, G., Yang, D., Wu, L., Shen, Q., et al.: Knowledge, attitudes and practices concerning self-medication with antibiotics among university students in western China. *Trop. Med. Int. Health* **19**(7), 769–779 (2014). <https://doi.org/10.1111/tmi.12322>
89. Mihretie, T.M.: Self-Medication Practices with Antibiotics Among Urban Dwellers of Bahir Dar Town, North West Ethiopia. Addis Ababa University, Addis Ababa (2014)
90. Shah, S.J., Ahmad, H., Rehan, R.B., Najeel, S., Mumtaz, M., Jilani, M.H., et al.: Self-medication with antibiotics among non-medical university students of Karachi: a cross-sectional study. *BMC Pharmacol. Toxicol.* **15**, 74 (2014). <https://doi.org/10.1186/2050-6511-15-74>
91. Widayati, A., Suryawati, S., de Crespigny, C., Hiller, J.E.: Self medication with antibiotics in Yogyakarta City Indonesia: a cross sectional population-based survey. *BMC Res. Notes* **4**, 491 (2011). <https://doi.org/10.1186/1756-0500-4-491>
92. Hadi, U., Duerink, D.O., Lestari, E.S., Nagelkerke, N.J., Werter, S., Keuter, M., et al.: Survey of antibiotic use of individuals visiting public healthcare facilities in Indonesia. *Int. J. Infect. Dis.* **12**(6), 622–629 (2008). <https://doi.org/10.1016/j.ijid.2008.01.002>
93. Al-Azzam, S.I., Al-Husein, B.A., Alzoubi, F., Masadeh, M.M., Al-Horani, M.A.: Self-medication with antibiotics in Jordanian population. *Int. J. Occup. Med. Environ. Health* **20**(4), 373–380 (2007). <https://doi.org/10.2478/v10001-007-0038-9>
94. Sawair, F.A., Baqain, Z.H., Abu Karaky, A., Abu, E.R.: Assessment of self-medication of antibiotics in a Jordanian population. *Med. Princ. Pract.* **18**(1), 21–25 (2009). <https://doi.org/10.1159/000163041>
95. Awad, A., Eltayeb, I., Matowe, L., Thalib, L.: Self-medication with antibiotics and antimalarials in the community of Khartoum State, Sudan. *J. Pharm. Pharm. Sci.* **8**(2), 326–331 (2005)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com